

Measles Measles is an infection that spreads very easily and can cause serious problems in some people. Having the MMR vaccine is the best way to prevent it.

Check if you or your child has measles

It has been confirmed that a child in Year 1 Violet has measles. If your child has had the MMR vaccine, it is unlikely that they will get it but they are more at risk if they have not.

Measles usually starts with cold-like symptoms, followed by a rash a few days later. Some people may also get small spots in their mouth.

Cold-like symptoms

The first symptoms of measles include:

- a high temperature
- a runny or blocked nose
- sneezing
- a cough
- red, sore, watery eyes

Spots in the mouth



Small white spots may appear inside the cheeks and on the back of the lips a few days later. These spots usually last a few days.

The measles rash

A rash usually appears a few days after the cold-like symptoms.



The rash starts on the face and behind the ears before spreading to the rest of the body.



The spots of the measles rash are sometimes raised and join together to form blotchy patches. They're not usually itchy.



The rash looks brown or red on white skin. It may be harder to see on brown and black skin.

Information:

If you're not sure it's measles

It's very unlikely to be measles if you've had both doses of the MMR vaccine or you've had measles before.

Urgent advice: Ask for an urgent GP appointment or get help from NHS 111 if:

- you think you or your child may have measles
- you've been in close contact with someone who has measles and you've not had measles before or you've not had 2 doses of the MMR vaccine
- you've been in close contact with someone who has measles and you're pregnant – measles can be serious in pregnancy
- you have a weakened immune system and think you have measles or have been in close contact with someone with measles
- Measles can spread to others easily. Call your GP surgery before you go in. They may suggest talking over the phone.
- You can also call 111 to get advice

How to look after yourself or your child

Measles usually starts to get better in about a week.

After seeing a GP, there are things you can do to help ease the symptoms and reduce the risk of spreading the infection.

It can help to:

- rest and drink plenty fluids, such as water, to avoid dehydration
- take paracetamol or ibuprofen to relieve a high temperature – **do not** give aspirin to children under 16 years
- use cotton wool soaked in warm water to gently remove any crusts from your or your child's eyes

Important

Stay off nursery, school or work for at least 4 days from when the rash first appears.

Also try to avoid close contact with babies and anyone who is pregnant or has a weakened immune system.

How to avoid spreading or catching measles

Measles is spread when an infected person coughs or sneezes. There are things you can do to reduce the risk of spreading or catching it.

Do

- wash your hands often with soap and warm water
- use tissues when you cough or sneeze
- throw used tissues in the bin

Don't

- do not share cutlery, cups, towels, clothes, or bedding